Sep.17, 2010

CURRICULUM VITAE
For
Osamu Tamada

Work Address:

Faculty of Bioenvironmental Science, Kyotogakuen University

Sogabe-cho, Kameoka, Kyoto 621-8555, Japan

Work Telephone:

81-771-29-3468

E-mail:

tamada@t02.mbox.media.kyoto-u.ac.jp

Home Address:

Nanmei-cho 13-36, Fukakusa, Fushimi-ku, Kyoto 612-0802, Japan

Home Telephone:

81-75-551-9126

Place of Birth:

Kyoto City

Citizenship:

Japan

Date of Birth:

Feb.16, 1944

EDUCATION

Kyoto University	1968	B.S. Faculty of Science, Department of Physics
Kyoto University	1970	Research Trainee, Geology and Mineralogy
Kyoto University	1973	M.S. Faculty of Science, Department of Geology and Mineralogy
Kyoto University	1980	Ph.D. Faculty of Science, Department of Geology and Mineralogy

EMPLOYMENT

Matsushita Electric Industrial Co.Ltd	1968	Technical Center
Kyoto University	1974	Assistant professor, Department of Liberal Arts and Sciences
Kyoto University	1980	Associate Professor, Department of

		Liberal Arts and Sciences
Kyoto University	1992	Professor, Graduate School of Human and Environmental Studies
Kyotogakuen University	2007	Professor, Department of Bioenvironmental Science

PROFESSIONAL MEMBERSHIPS

Mineralogical Society of Japan	1972-present
Mineralogical Society of America	1978-present
Crystallographic Society of Japan	1986-2007

JOURNAL PUBLICATIONS (AFTER 2000)

Kitashita, K., Hagiwara, R., Ito, Y. and **Tamada, O**. (2000) Crystal structure of some cubic hexafluorophosphates at ambient temperatures. J. Fluorine Chem., 101, 173-179.

Hagiwara, R., Kitashita, K., Ito, Y. and **Tamada, O.** (2000) Crystal structure of $AgPF_6$ and $AgAsF_6$ at ambient temperatures. Solid State Sciences, 2, 237-241.

Hori, T., Taniguti N., Oozono, T., **Tamada, O.** (2001) Actual method displaying c rystal structure using a devise to present 3-dimensional coordinates (in Japanese). Chemistry and Education, 49, 730-733.

Matsumoto, K., Hagiwara, R., Ito, Y. and **Tamada, O.** (2001) Crystal structure of $AgAF_6$ (A=P, As, Sb, Nb, Ta) at ambient temperatures. J. Fluorine Chem., 110, 117-122.

Matsumoto, K., Tsuda, T., Nohira,T., Hagiwara, R., Ito, Y. and **Tamada, O.** (2002) Tris (1-ethyl-3-methylimidazolium) hexa-chlorolanthanate. Acta Cryst., C58, m186-m187.

Matsumoto, K., Tsuda, T., Hagiwara, R., Ito,Y. and **Tamada, O.** (2002) Structural characteristics of 1-ethyl-3-methylimidazolium bifluoride : HF-deficient form of a highly conductive room temperature molten salt. Solid State Sciences, 4, 23-26.

Ito, K., Sato, H., Takei, H., **Tamada, O.** and Kitazawa, T. (2002) Synthesis of la rge high-quality forsterite single crystals to 200 mm length and its significance. Geochem. Geophys. Geosystems, 4, 1011(1-4).

Matsumoto, K., Hagiwara, R., Ito, Y. and **Tamada, O.** (2002) Synthetic, structur al and thermal studies of Ag(I)-XeF_2 complex salts. Solid State Sciences, 4, 1465-1469.

Ito, K., Sato, H., Kanazawa, H., Kawame, N., **Tamada, O.**, Miyazaki, K., Uehara, S., Iio, Y., Takei, K., Kitazawa, T., Koike, M., Matsushita, Y. and Ito, Y. (2003) First synthesis of olivine single crystal as large as 250 carats. J. Cryst. Growth, 253, 557-561.

Ikuma, N., Tamura, R., Shimono, S., Kawame, N., **Tamada, O.**, Sakai, N., Yamauchi, J. and Yamamoto,Y. (2003) Crystal structure and magnetic properties of novel chiral nitroxides existing as racemic conglomerates. Mendeleav Commun., 109-111.

Shimono, S., Tamura, R., Ikuma, N., Takimoto, T., Kawame, N., **Tamada, O.**, Sakai, N., Matsuura, H. and Yamauchi, J. (2004) Preparation and characterization of new chiral nitronyl nitroxides bearing a stereogenic center in the imidazolyl framework. J. Org. Chem., 69, 475-481.

Ikuma, N., Tamura, R., Shimono, S., Kawame, N., **Tamada, O.**, Sakai, N.,Yamauchi, J. and Yamamoto, Y. (2004) Magnetic properties of all-organic liquid crystals containing a chiral five-membered cyclic nitroxide unit within the rigid core. Angew. Chem. Int. Ed., 43, 3677-3682.

Ito, K., Sato, H., Kanazawa, H., **Tamada, O.**, Kitazawa, T., Koike, M. and Takei, H. (2004) Simple practical processing of a large single crystal from a large amou nt of a complete stoichiometric mixture in two-component system. J. Cryst. Growt h, 271, 398-402.

Wajima, T., Kuzawa, K., Ishimoto, H., **Tamada, O.**, and Nishiyama, T. (2004) T he synthesis of zeolite-P, Linde Type A, and hydroxysodalite zeolites from paper s ludge ash at low temperature (80°C): Optimal ash-leaching condition for zeolite sy nthesis. Am. Mineral., 89, 1694-1700.

Ikuma, N., Tamura, R., Shimono, S., Kawame, N., **Tamada, O.**, Sakai, N., Yamamoto, Y., Yamauchi, J. (2005) Characterization of the crystalline nature of the racemates of novel chiral five-membered cyclic nitroxides. Molecular Crystals and Liquid Crystals, 440, 23-35.

Wajima, T., Haga, M., Kuzawa, K., Ishimoto, H., **Tamada, O.**, Ito, K., Nishiyama, T. (2006) Zeolite synthesis from paper sludge ash at low temperature (90 °C) with addition of diatomite. J. Hazardous Materials, 132, 244－252.

Rakovan, J., Kitamura, M., **Tamada, O.** (2006) Sakura ishi (cherry blossom stones) : Mica pseudomorphs of complex cordierite-indialite intergrowths from Kameoka, Kyoto Prefecture, Japan. Rocks & Minerals, 81, 284－292.

Ikuta, D., Kawame, N., Banno, S., Hirajima, T., Ito, K., Rakovan, J. F., Downs, R. T., **Tamada, O.** (2007) First in situ X-ray identification of coesite and retrograde quartz on a glass thin section of an ultrahigh-pressure metamorphic rock and their crystal structure details. Am. Mineral., 92, 57－63.

Kawame, N., Ikuta, D., Kanazawa, H., Ito, K., Gunter, E. M., Boisen, B. M. and **Tamada, O.** (2007) Superstructure of Challis mordenite with doubled monoclinic unit cell. Am. Mineral., 92, 892-897.

Kanazawa, H., Ito, K., Sato, H., Kumatoriya, M., Miyazaki, K., Uehara, S., Tsuda, H., Kuzawa, K., Kawame, N., Kitazawa, T., **Tamada, O.**, Boisen, M. B., Takei, H. (2007) Synthesis and absorption spectra of large homogeneous single crystals of forsterite doped with manganese. J. Cryst. Growth, 304, 492-496.

Wajima, T., Ishimoto, H., Kuzawa, K., Ito, K., **Tamada, O.**, Gunter, M. E., Rakovan, J. F. (2007) Material conversion from paper-sludge ash in NaOH, KOH and LiOH solutions. Am. Mineral., 92, 1105－1111.

Nakashima, K., Kawame, N., Kawamura, Y., **Tamada, O.** and Yamauchi, J. (2009) Bis (2, 3, 5-triphenyltetrazolium) tetrathiocyanatocobaltate (II). Acta Cryst., E65, m1406-m1407.

BOOKS (AFTER 2000)

Tamada, O. (2000) Can we continue to consume natural resources and energy ? (in Japanese). In "Paradigm Shift in 20th Century" edited by M. Hirai, Kokusyo Kankoukai, Chapter 3, 69-87.

Gibbs, G. V., Boisen, M. B., Hill, F. C.and **Tamada, O.** (2000) Search for a connection among bond strength, bond length and electron density distributions. In "Physics Meets Mineralogy --- Condensed-Matter Physics in Geosciences ---". edited by H. Aoki, Y. Syono and R. J. Hemley, Cambridge University Press Part III, Chapter 3.1, 83-94.