



University of Idaho Extension

Jim Wasia reported the first 2018 incidence of **stripe rust in the Aberdeen** vicinity, and I am buying him lunch.

Literally three minutes after, Brad Clayson, CHS in Blackfoot sent a picture of a hot spot of stripe rust in Brundage south of Idaho Falls.

Overall, stripe rust pressure is low and showing in the susceptible winter wheat. There should be very little loss in winter wheat. We will keep an eye on the development in spring wheat, but be sure to be vigilant in scouting susceptible and moderately susceptible spring wheat varieties. There are tables in the back of the 2017 Small Grains Research Report at our website <http://www.uidaho.edu/extension/cereals/scseidaho>

The entire report can be downloaded as a PDF
at <http://www.extension.uidaho.edu/publishing/pdf/RES/RES193.pdf>
The stripe rust results for spring wheat is on Addendum 5 page 135.

For fungicide efficacy please refer to the attached chart of the NCERA-184 Wheat Fungicide Efficacy Chart for 2018. Resistant and moderately resistant varieties **SHOULD NOT** need fungicide application to control stripe rust. Susceptible or moderately susceptible varieties of winter wheat and spring wheat should be sprayed when 5% incidence occurs in the field.

Keep reporting!

cheers,
Juliet

Management of Small Grain Diseases Fungicide Efficacy for Control of Wheat Diseases (2018 Final)

The North Central Regional Committee on Management of Small Grain Diseases (NCERA-184) has developed the following information on fungicide efficacy for control of certain foliar diseases of wheat for use by the grain production industry in the U.S. Efficacy ratings for each fungicide listed in the table were determined by field testing the materials over multiple years and locations by the members of the committee. Efficacy is based on proper application timing to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are based on a single application of the labeled rate as listed in the table. Table includes most widely marketed products, and is not intended to be a list of all labeled products.

Efficacy of fungicides for wheat disease control based on appropriate application timing

Fungicide(s)				Powdery mildew	Stagonospora leaf/glume blotch	Septoria leaf blotch	Tan spot	Stripe rust	Leaf rust	Stem rust	Head scab ⁴	Harvest Restriction
Class	Active ingredient	Product	Rate/A (fl. oz)									
Strobilurin	Picoxystrobin 22.5%	Aproach SC	6.0 – 12.0	G ¹	VG	VG ²	VG	E ³	VG	VG	NL	Feekes 10.5
	Fluoxastrobin 40.3%	Evito 480 SC	2.0 – 4.0	G	--	--	VG	--	VG	--	NL	Feekes 10.5 and 40 days
	Pyraclostrobin 23.6%	Headline SC	6.0 - 9.0	G	VG	VG ²	E	E ³	E	G	NL	Feekes 10.5
Triazole	Metconazole 8.6%	Caramba 0.75 SL	10.0 - 17.0	VG	VG	--	VG	E	E	E	G	30 days
	Tebuconazole 38.7%	Folicur 3.6 F ⁵	4.0	NL	NL	NL	NL	E	E	E	F	30 days
	Prothioconazole 41%	Proline 480 SC	5.0 - 5.7	--	VG	VG	VG	VG	VG	VG	G	30 days
	Prothioconazole 19% Tebuconazole 19%	Prosaro 421 SC	6.5 - 8.2	G	VG	VG	VG	E	E	E	G	30 days
	Propiconazole 41.8%	Tilt 3.6 EC ⁴⁵	4.0	VG	VG	VG	VG	VG	VG	VG	P	Feekes 10.5.4
Mixed modes of action ⁶	Tebuconazole 22.6% Trifloxystrobin 22.6%	Absolute Maxx SC	5.0	G	VG	VG	VG	VG	E	VG	NL	35 days
	Cyproconazole 7.17% Picoxystrobin 17.94%	Aproach Prima SC	3.4 - 6.8	VG	VG	VG	VG	E	VG	--	NR	45 days
	Prothioconazole 16.0% Trifloxystrobin 13.7%	Delaro 325 SC	8.0	G	VG	VG	VG	VG	VG	VG	NL	Feekes 10.5 35 days
	Fluapyroxad 2.8% Pyraclostrobin 18.7% Propiconazole 11.7%	Nexicor EC	7.0 - 13.0	G	VG	VG	E	E	E	VG	NL	Feekes 10.5
	Fluoxastrobin 14.8% Flutriafol 19.3%	Preemptor SC	4.0 - 6.0	--	--	VG	VG	E	VG	--	NL	Feekes 10.5 and 40 days
	Fluxapyroxad 14.3% Pyraclostrobin 28.6%	Priaxor	4.0 - 8.0	G	VG	VG	E	VG	VG	G	NL	Feekes 10.5
	Propiconazole 11.7% Azoxystrobin 13.5%	Quilt Xcel 2.2 SE ⁵	10.5 - 14.0	VG	VG	VG	VG	E	E	VG	NL	Feekes 10.5.4
	Prothioconazole 10.8% Trifloxystrobin 32.3%	Stratego YLD	4.0	G	VG	VG	VG	VG	VG	VG	NL	Feekes 10.5 35 days
	Benzovindiflupyr 2.9% Propiconazole 11.9% Azoxystrobin 10.5%	Trivapro SE	9.4 - 13.7	VG	VG	VG	VG	E	E	VG	NL	Feekes 10.5.4 14 days

¹Efficacy categories: NL=Not Labeled; NR=Not Recommended; P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent; -- = Insufficient data to make statement about efficacy of this product.

² Product efficacy may be reduced in areas with fungal populations that are resistant to strobilurin fungicides.

³Efficacy may be significantly reduced if solo strobilurin products are applied after stripe rust infection has occurred.

⁴Application of products containing strobilurin fungicides may result in elevated levels of the mycotoxin Deoxynivalenol (DON) in grain damaged by head scab.

⁵Multiple generic products containing the same active ingredients also may be labeled in some states.

⁶Products with mixed modes of action generally combine triazole and strobilurin active ingredients. Nexicor, Priaxor and Trivapro include carboxamide active ingredients.