201100051

## THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

## University of Idaho

Whereas, there has been presented to the

## Secretary of Agriculture

An application requesting a certificate of protection for an alleged distinct variety of sexually reproduced, or tuber propagated plant, the name and description of which are contained in the application and exhibits, a copy of which is hereunto annexed and made a part hereof, and the various requirements of LAW in such cases made and provided have been complied with, and the title thereto is, from the records of the PLANT VARIETY PROTECTION OFFICE, in the applicant(s) indicated in the said copy, and Whereas, upon due examination made, the said applicant(s) is (are) adjudged to be entitled to a certificate of plant variety protection under the LAW.

Now, therefore, this certificate of plant variety protection is to grant unto the said applicant(s) and the successors, heirs or assigns of the said applicant(s) for the term of TWENTY years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to exclude others from selling the variety, or offering it for sale, or reproducing it, or importing it, or exporting it, or conditioning it for propagation, or stocking it for any of the above purposes, or using it in producing a hybrid or different variety therefrom, to the extent provided by the PLANT VARIETY PROTECTION ACT. IN THE UNITED STATES SEED OF THIS VARIETY (1) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)



Attest:

02-3

Commissioner

RAPE

'Clearwater'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C. this twenty-fifth day of June, in the year two thousand and fourteen.

Clean J. Vilval

Secretary of Agriculture

	ate on all reproductions		Form Approved - OMB No. 0581-00
AGRICULTURAL M	T OF AGRICULTURE ARKETING SERVICE ANT VARIETY PROTECTION OFFICE	The following statements are made in accordant the Paperwork Reduction Act (PRA) of 1995.	ce with the Privacy Act of 1974 (5 U.S.C. 552a) and
APPLICATION FOR PLANT VAR	ELETY PROTECTION CERTIFICATE action burden statement on reverse)	Application is required in order to determine if a (7 U.S.C. 2421). Information is held confidential	plant variety protection certificate is to be issued until certificate is issued (7 U.S.C. 2426).
1. NAME OF OWNER	on burden statement on reverse)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NAME	3. VARIETY NAME
University of Idaho		IMI.SUN.56/UI.C.1.04	Clearwater
4. ADDRESS (Street and No., or R.F.D. No., Ci	ty, State, and ZIP Code, and Country)	5. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY
Office of Technology Tran	sfer 875 Perimeter Driv	ve,(208) 885 4550	
PO Box 443003 Morill Hall 414 Morrill }	[-11-414 MS3003	6. FAX (include area code)	┥
Moscow, Idaho, 83844-30	03	(208) 885 4551	#20110005
7. IF THE OWNER NAMED IS NOT A "PERSO FORM OF ORGANIZATION (corporation, partnerssociation, etc.)	E OWNER NAMED IS NOT A "PERSON", GIVE 8. IF INCORPORATED, GIVE 9. DATE OF INCORPORATION  OF ORGANIZATION (corporation, partnership, STATE OF INCORPORATION		
University of Idaho			
-Jack Brown Inclu-PSES, CALS Karen S	meter Drive, MS30003		C CERTIFICATE FEE:  S V E DATE
11. TELEPHONE (Include area code)	12. FAX (Include area code)	13. E-MAIL	D
(208) 885 7078	(208) 885 7760	karens ibrown@uidaho.edu	1 & gaylene@uidaho.edu=
14. CRÓP KIND (Common Name)  Spring (canola)  15. GENUS AND SPECIES NAME OF CROP	16. FAMILY NAME (Botanical) Brassicaceae	☐ YES X NO IF SO, PLEASE GIVE THE ASSIGN	ED USDA-APHIS REFERENCE NUMBER FOR THE
15. GENUS AND SPECIES NAME OF CROP  Brassica napus L.	17. IS THE VARIETY A FIRST GENERATION I	HYBRID? APPROVED PETITION TO DEREG	GULATE THE GENETICALLY MODIFIED PLANT FOR
19. CHECK APPROPRIATE BOX FOR EACH A (Follow instructions on reverse)		20. DOES THE OWNER SPECIFY	THAT SEED OF THIS VARIETY BE SOLD ONLY AS
a. X Exhibit A. Origin and Breeding History of the	ne Variety		See Section 83(a) of the Plant Variety Protection Act) and 22 below) □ NO (If "no", go to item 23) □
b. X Exhibit B. Statement of Distinctness		UNDECIDED	
c. X Exhibit C. Objective Description of Variety		21. DOES THE OWNER SPECIFY	THAT SEED OF THIS VARIETY BE LIMITED AS TO
d. X Exhibit D. Additional Description of the Var	iety (Optional)	NUMBER OF CLASSES?  X YES □ NO	
e. X Exhibit E. Statement of the Basis of the Ov	vner's Ownership		UNDATION □ REGISTERED X CERTIFIED
f. X Exhibit F. Declaration Regarding Deposit			THAT SEED OF THIS VARIETY BE LIMITED AS TO
g. X Voucher Sample (3,000 viable untreated at that tissue culture will be deposited and maintain	eeds or, for tuber propagated varieties, verification	NUMBER OF GENERATIONS?	THE SELECTION PARTIES AS TO
h. K Filing and Examination Fee (\$4,382), mad	e payable to "Treasurer of the United	☐ YES X NO IF YES, SPECIFY THE NUMBER 1,	2,3, etc. FOR EACH CLASS.
States" (Mail to the Plant Variety Protection Office Pages remitted per cc pymat	e) • (tabe u /valesessa)	☐FOUNDATION ☐REGISTER	ED CERTIFIED
23 HAS THE VARIETY (INCLUDING ANY HAR	VESTED MATERIALLOR A HYPRID PRODUCTI		ry, please use the space indicated on the reverse.) IPONENT OF THE VARIETY PROTECTED BY
FROM THIS VARIETY BEEN SOLD, DISPOSED OTHER COUNTRIES?	OF, TRANSFERRED, OR USED IN THE U. S. (		T (PLANT BREEDER'S RIGHT OR PATENT)?
FOR EACH COUNTRY AND THE CIRCUMSTAI	IRST SALE, DISPOSITION, TRANSFER, OR US NCES. (Please as space indicated on reverse.)	E IF YES, PLEASE GIVE COUNTRY, REFERENCE NUMBER. (Please us	DATE OF FILING OR ISSUANCE AND ASSIGNED be space indicated on reverse.)
	I be deposited in a public repository and maintain		ccordance with such regulations as may be applicable,
	is sevinally reproduced or tubor proposed alert		t, uniform, and stable as required in Section 42, and is
Owner(s) is (are) informed that false representati	on herein can jeopardize protection and result in		
by Dow		SIGNATURE OF OWNER  A  A  A  A  A  A  A  A  A  A  A  A  A	h
NAME (Please drint or type)		NAME (Please print or type)	
Jack Brown	DATE	GAYLENE A	UDERSON

2010 MOV 10 FH 5:18

GENERAL INSTRUCTIONS: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E, F; (3) for a tuber reproduced variety, verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; and (4) payment by credit card or check drawn on a U.S. bank for \$4,382 (\$518 filing fee and \$3,864 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice). NEW: With the application for a seed reproduced variety or by direct deposit soon after filing, the applicant must provide at least 3,000 viable untreated seeds of the variety per se, and for a hybrid variety at least 3,000 untreated seeds of each line necessary to reproduce the variety. Partial applications will be held in the PVPO for not more than 90 days; then returned to the applicant as un-filed. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a payment by credit card or check payable to "Treasurer of the United States" in the amount of \$768 for issuance of the certificates will be issued to owner, not licensee or agent.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filling a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

Plant Variety Protection Office

Telephone: (301) 504-5518 FAX: (301) 504-5291

General E-mail: PVPOmail@usda.gov

Homepage: http://www.ams.usda.gov/science/pvpo/PVPindex.htm

#### SPECIFIC INSTRUCTIONS:

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that the permanent name of the application variety (even if it is a parental, inbred line) has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: U.S. Department of Agriculture, Agricultural Marketing Service, Livestock and Seed Programs, Seed Regulatory and Testing Branch, 801 Summit Crossing Place, Suite C, Gastonia, North Carolina 28054-2193 Telephone: (704) 810-8870. http://www.arns.usda.gov/lsg/seed.htm.

#### ITEM

- 19a. Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
  - (2) the details of subsequent stages of selection and multiplication;
  - (3) evidence of uniformity and stability; and
  - (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the
- new variety is most similar to one variety or a group of related varieties:
  - (1) identify these varieties and state all differences objectively;
  - (2) attach replicated statistical data for characters expressed numerically and demonstrate that these are clear differences; and
  - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)
- 24. CONTINUED FROM FRONT (Please give the country, date of filling or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

## 'Clearwater' Spring Rapeseed Brassica napus L.

### **Exhibit A: Origin and Breeding History**

'Clearwater' spring rapeseed (*Brassica napus* L. spp. *oleifera* (Metzg) Sinsk. *f. annua*) was developed for use as an edible oil-quality (canola) cultivar by the Idaho Agricultural Experimental Station, Moscow, ID 83844. This cultivar is protected by U.S. Plant Variety Protection (PVP pending).

Clearwater is a near pure-line spring canola cultivar which has high resistance to imidazolinone class herbicides including imazamox (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid). Clearwater produces canola-quality seed oil and canola-quality seed meal, and was selected for adaptability to environments throughout the Pacific Northwest Region (Idaho, Oregon, Washington, and Montana). This cultivar was developed from the cross 'CFSN.39.9'/ 'Sunrise'////'Sunrise'. CFSN.39.9 is a canola-quality breeding selection with low erucic acid content (less than 20 g kg<sup>-1</sup>) and less than 30  $\mu$ mol g<sup>-1</sup> of glucosinolate in the defatted seed meal, which was selected from a segregating  $F_3$  population derived from a cross 'Profit'/'Springfield'. Progeny from this cross were accidentally planted into soil with a high imazamox residue and CFSN.39.9 was one of 55 single plants that survived from a total of 5 acres of segregating breeding lines planted. CFSN.39.9 was later shown to have the  $PM_1$  and  $PM_2$  mutation genes conferring tolerance to imidazolinone class herbicides. Sunrise is a canola-quality spring rapeseed cultivar developed and released by the University of Idaho in 1997 (Brown et al., 1997).

Seed from the original cross CFSN.39.9/Sunrise was produced and  $F_1$  progeny was planted in the glasshouse and backcrossed to Sunrise to produce  $BC_1F_1$  seed. Fifty seeds of the  $BC_1F_1$  seed were planted in seedling trays and grown in the glasshouse. When plants reached the 4-6 leaf stage (approximately three weeks later), they were sprayed with imazamox herbicide at a rate equivalent to 35 g a.i. ha<sup>-1</sup>, along with non-ionic surfactant 2.5 ml l<sup>-1</sup> of solution liquid urea-ammonium sulfate fertilizer 15 ml l<sup>-1</sup> of solution. Seedlings that showed no visible herbicide damage symptoms were transplanted to larger pots and these plants were backcrossed to Sunrise to produce  $BC_2F_1$  seed. The procedure of spraying seedlings with imazamox herbicide, identifying highly tolerant seedlings and using these to backcross to Sunrise was repeated a further three times to produce  $BC_5F_1$  seed. Five hundred  $BC_5F_1$  seed were planted into seedling trays and sprayed with imazamox herbicide at the above rate. Seedlings expressing high tolerance (156 lines) to the herbicide spray were selected and grown to maturity in large pots to produce  $BC_5F_2$  seed.

 $BC_5F_2$  seed harvested from the glasshouse was used to plant unreplicated single plant plots (15 m x 2 rows 15 cm apart) in the field.

Single plant plots were sprayed at the 4 leaf stage with imazamox herbicide at described

above, and evaluated for herbicide tolerance along with morphological adaptability characters, seedling vigor, flower date, plant height, lodging and maturity. At maturity, seed from 26 single plant plots were selected for further evaluation. Ten single plants were harvested separately from each plot while the remainder of the plots was hand threshed and used for planting yield trials. The single BC<sub>5</sub>F<sub>3</sub> plants harvested were planted as single plant plots and used to initiate seed increases.

Twenty six BC<sub>5</sub>F<sub>3</sub> selected lines were evaluated in replicated yield trials at two locations in northern Idaho. Included in the trial were several commercially available spring canola cultivars. None of the yield evaluation trials were sprayed post emergence with imazamox herbicide. A range of morphological characters was recorded including seedling vigor, flower date, plant height, lodging and maturity. After harvest, a sample of seed was taken from each plot to determine oil content, fatty acid profile (Hammond, 1991; Christie, 1992) and glucosinolate content was estimated using a Tes-tape<sup>®</sup> procedure (Smith & Donald, 1988). After reviewing the performance of the 26 BC<sub>5</sub>F<sub>3</sub> selections in yield trials and as single plant plots, the line code IMI.SUN.56 was selected, and 200 single plant selections from this BC<sub>5</sub>F<sub>4</sub> family (later coded as UI.C.1) were harvested used to initial the seed source for the Clearwater cultivar.

A single seed from each of the 200 BC<sub>5</sub>F<sub>4</sub> single plant selections from UI.C.1 was planted in the glasshouse. At the seedling stage, a leaf DNA extract was taken from each seedling and screened for the presence of both tolerance genes ( $PM_1$  and  $PM_2$ ). Seedlings that were homozygous for both tolerance alleles were grown to maturity, and the resulting seed was planted to initiate a pure seed source for increase in the field. At each stage of seed increase seedlings were sprayed at the 4 leaf stage with imazamox herbicide as described above, and plots showing any susceptibility symptoms were discarded. Breeders' seed (UI.C.1.04) was produced in 2004 from 400 single plant selections made from the BC<sub>5</sub>F<sub>7</sub> family. Foundation seed was produced from Breeders seed in 2005. Throughout the stages of Clearwater seed increases including pre-Breeders seed, Breeders seed, Foundation seed, and finally Certified seed production, plants were consistently uniform and stable, and no variants were observed over this four year period.

#### References

Brown, J., J.B. Davis, A.P. Brown, D.A. Erickson and L. Seip, 1997. Registration of 'Sunrise' spring rapeseed. *Crop Sci.* 38:542-543.

Christie, W.W., 1992. Preparation of fatty acid methyl ester. *Inform* 3:1031-1034.

Hammond, E.G., 1991. Organization of rapid analysis of lipids in many individual plants. *In*: Modern Methods of Plant Analysis. Vol: 12. Springer-Verlag, Berlin, Germany.

Smith, D.B., and C.N. Donald, 1988. The measurement of glucosinolates in oilseed rape by glucose reaction. *Plant Varieties and Seeds* 1:121-130.

## 'Clearwater' Spring Rapeseed Brassica napus L.

#### **Exhibit B: Statement of Distinctness**

Clearwater is most similar in plant appearance to the spring canola cultivar Sunrise (Brown et al., 1997). However, Clearwater is high tolerant/resistance to imidazolinone class herbicides including imazamox (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1*H*-imidazol-2-yl]-5-(methoxymethyl)-3-pyridinecarboxylic acid) while Sunrise is highly susceptible to all imidazolinone class herbicides including imazamox. In field trials conducted at Moscow and Genesee in 2004 and 2005 Clearwater was grown in replicated field plots alongside Sunrise and seedlings sprayed with a 1X rate of Beyond<sup>®</sup> (imazamox) herbicide. Stand counts were conducted on a single 1 m row from each plot before and after spraying and %survival data are presented in Table B1.

**Table B1.** Percentage of Clearwater and Sunrise seedling that survival after seedlings had been sprayed with Beyond (imazamox) herbicide.

1 2	•	,			
Cultivar	Average	2005		20	04
		Moscow	Genesee	Moscow	Genesee
		%	survival afte	er treatment -	
Clearwater	99	101	98	97	100
Sunrise	0	0	0	0	0

#### References

Brown, J., J.B. Davis, A.P. Brown, D.A. Erickson and L. Seip, 1997. Registration of 'Sunrise' spring rapeseed. *Crop Sci.* 38:542-543.

REPRODUCE LOCALLY. Include form number and date on all reproductions. Form Approved OMB NO 0581-0055

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (20 720-2600 (voice and TDD).

To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF EXHIBIT C
AGRICULTURAL MARKETING SERVICE
SCIENCE AND TECHNOLOGY
PLANT VARIETY PROTECTION OFFICE
BELTSVILLE, MD 20705

# OBJECTIVE DESCRIPTION OF VARIETY RAPESEED (Brassica napus and B. campestris)

NAME OF APPLICANT (S)	TEMPORARY OR EXPERIMENTAL DESIGNATION	VARIETY NAME
University of Idaho Agricultural	IMI.SUN.56/UI.C.1.04	Clearwater
Experiment-Station		
ADDRESS (Street and No. or RD No.	o., City, State, Zip Code, and Country)	FOR OFFICIAL USE ONLY
University of Idaho		PVPO NUMBER
Moscow, Idaho, 83844-2339		201100051
1. SPECIES		
* _X_ Brassica napus Brassic	ca campestris	
2. TYPE		
* _X_ Spring Winter		
3. PLANT HEIGHT (at pod mat	urity)	
	re to standard variety below)	
cm shorter than Chec	ck variety:	
Height same as Check variety: _		
89 cm taller than Che	ck variety: <u>Sunrise</u>	
* Height class: Autumn sown	_2_ Spring sown	
	1 = Short (Erglu)	
	2 = Medium short ()	
	3 = Medium (Cresus) 4 = Medium tall ( )	
	5 = Tall (Petranova)	
4. STEM ANTHOCYANIN		
1 1 Aboont 2 Wook 2 Mo	dium 4 Ctrong	
1 1 = Absent 2 = Weak 3 = Med 5. <b>SEED COTYLEDONS</b> (maxim	mum width fully developed; mean of 50 graded seeds)	
_2_1 = Narrow (Erglu) 2 = Mediu	m (Primor) 3 = Broad (Expander)	
6. SEEDLING GROWTH HABI	T (leaf rosette)	

\_1\_ 1 = Upright 2 = Prostrate (short photoperiod)

- \* 1 Anther dotting (at opening of flower; given percentage
- %) 1 = Absent (X) 2 = Few () 3 = Medium (Primor) 4 = Many ()
- Flowering class (Autumn sown) 4 Flowering class (Spring sown): 58 Days after planting to 50% flower bloom, same as 'Sunrise'

  1 = Very early (Arvor) 1 = Very early (Tower)

  2 = Early (Primor) 2 = Early (Kosa)

  3 = Medium early () 3 = Medium early ()

  4 = Medium late () 4 = Medium late (X)

  5 = Late (Marcus) 5 = Late (Petranova)

  6 = Very late () 6 = (Very late)

#### 9. PODS (Slique)

- \* <u>1</u> Pod type: 1 = Bilateral single pod (Jet Neuf) 2 = Other ()
- \* 2 Silique beak length: (given length: 8.91 mm)1 = Short (Forto) 2 = Medium (Liragold) 3 = Long (Rapol)
- \* \_2\_ Pod length; (give length: \_61\_.\_17 mm) 1 = Short ( ) 2 = Medium (X) 3 = Long ( )
- \* 3 Pod width; (give width: 5 91 mm) 1 = Narrow () 2 = Medium (X) 3 = Wide ()
- \* \_4\_ Pod habit: 1 = Erect (Gulliver) 2 = Semi-erect to erect (Oro) 3 = Semi-erect 4 = Horizonal to semi-erect (Brink) 5 = Horizonal
- \* 2 Pedicel length: (given length 18.34 mm) 1 = Very short () 2 = Short () 3 = Long ()
- \* \_4\_ Ripening Class (Autum sown): 1 = Very early () 2 = Early (X) 3 = Medium () 4 = Late (X) 5 = Very late ()
- \* \_1\_ \_1\_ \_7\_ Days to Maturity
- \* \_\_\_ \_ Days earlier than Check variety: \_\_\_\_\_
- \* Maturity same as Check variety: \_Sunrise
- \* \_\_\_\_\_\_7\_ Days later than Check variety: \_Hyola.401

#### 10. SEEDS

- \* <u>3</u>.<u>4</u> g/1000 unsized seed
- \* \_\_\_\_ g less than Check variety: \_\_\_\_\_
- \* Weight same as Check variety: \_Sunrise
- \* <u>0</u>.<u>4</u> g more than Check variety: <u>Hyola.401</u>
- \*  $\underline{2}$  Weight Class (grams): 1 = less than 3.0 (Candle) 2 = 3.0 3.9 (X) 3 = 4.0 5.0 (Jet Neuf) 4 = more than 5.0 ()
- $\underline{1-2}$  Seeds Per Pod: (give number:  $\underline{26.54}$  per pod): 1 = Low (X) 2 = Medium (X) 3 = High ()
- \* \_5\_ Testa Color: 1 = Black (Jet Neuf) 2 = Red ()
  3 = Yellow (Yellow Sarson) 4 = Dark to black (X)
  5 = Reddish-brown to black () 6 = Other \_\_\_\_\_

#### 11. CHEMICAL COMPOSITION OF SEED

- \* \_1\_ Euric Acid: 1 = Low (less than 2%) 2 = Intermediate 3 = High (more than 50%)
- \* \_1\_ Glucosinate Content; (give: \_\_\_ 9\_ .1\_ µmol/gram of defatted seed meal)
  1 = Low less than 30 µmol/gram of defatted seed meal (Candle) 2 = High More than 30 µmol/gram of defatted seed meal (Mikado)
- \* <u>39</u>.<u>1</u> % Oil

\_\_\_.\_\_ % Protein (oil free meal)

Fatty Acid Composition (%):

Palmitic	Stearic	Oleic	Linoleic	Linolenic	Eicosenoic	Erucic
16:0	18:0	18:1	18:2	18:3	20:1	22:1
* <u>4.1</u>	<u>2.2</u>	<u>62.1</u>	<u>19.5</u>	<u>7.6</u>	<u>1.4</u>	0.7

#### 12. FROST TOLERANCE (Late spring frosts)

\* 3 Tolerance: 1 = Not hardy - susceptible (Indore) 2 = Moderately suscesptible () 3 = Moderately resistant (X) 4 = Hardy (Bridger)

#### 13. LODGING RESISTANCE

\* 3 Resistance: 1 = Weak (Span) 2 = Moderately weak (Olga) 3 = Moderately strong (X) 4 = Strong (Torpe)

#### 14. HERBICIDE RESISTANCE

- \* <u>1</u> Atrazine: 1 = Susceptible (Jet Neuf) 2 = Resistant ()
- \* \_5\_ Imazamox (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1*H*-imidazol-2-yl]-50(methoxymethyl)-3-pyridinecarboxyl acid): 1 = Suscept (); 5 = Highly tolerant (X).
- 15. **DISEASE RESISTANCE** (0 = Not tested 1 = Susceptible 2 = Low resistance 3 = Moderate resistance 4 = High resistance)
- \* <u>0</u> Selerotinia Stem Rot (Scerotinia sclerotiorum)
- \* \_0\_ Black Let, Stem Canker (Leptosphaeria maculans, Plenodomus lingum, Phoma lingam)
- \* \_0\_ White Rust (Albugo candida, A. Cruciferrarum)
- \* <u>0</u> Light Leaf Spot (*Pyrenopeziza brassicae*)
- \* <u>0</u> Downy Mildew (*Peronospora parasitica*)
- \* <u>0</u> Rhizoctonia Root Rot (*Rhizoctonia solani*)
- \* \_0\_ Alternaria Black Spot (Alternaria brassicicola)
- \* \_0\_ Other \_\_\_\_\_

#### 16. COMMENTS (Please give any additional comments which characterizes the variety)

Glucosinolate composition of seed meal (µmol g-1 defatted seed meal

Destant	OLLD	Denterral	OLL Devices of
Butenyl	OH Butenyl	Pentenyl	OH Pentenyl
2.90	5.75	0.48	Trace

#### 17. DIRECTIONS

Select the number which characterizes the variety in the features above. **14**. Those characteristics marked with an asterisk "\*" should be recorded. Any others should be recorded if possible to help establish novelty or uniqueness. Characteristics described, including numerical measurements, should represent those that are <u>typical</u> for the variety. Give test area <u>Inland Pacific Northwest</u> conditions <u>2001 to 2007</u>.



## 'Clearwater' Spring Rapeseed Brassica napus L.

### Exhibit D: Additional Description of Variety

Clearwater leaves have medium serration of the leaf margins, medium lobing and no clasping of leaves attached to the stem (Figure D1). Clearwater plants flower on average 58.8 days after planting, which is not significantly different from Sunrise but significantly later than the early flowering cultivar Hyola.401 (Table D1). Clearwater plants are significantly taller (127.8 cm) than Sunrise (118.9 cm) and Hyola.401 (106.7 cm). Clearwater plants reached full maturity on average 97 days after planting, which was not significantly different from Sunrise which reaches maturity, on average, 96 days after planting. Average 1000-seed weight of Clearwater is 3.4 g, which is similar to Sunrise at 3.3 g. Clearwater is resistant to lodging and slightly resistant to seed shatter at maturity.

Clearwater was grown in replicated yield trials alongside Sunrise and Hyola.401 at a total of 79 year-sites between 2001 and 2007. None of these yield evaluation trial had imazamox herbicide applied. Averaged over all years and sites, Clearwater seed yield was 1,748 kg ha<sup>-1</sup>, compared to 1,646 kg ha<sup>-1</sup> for Sunrise and 1,986 kg ha<sup>-1</sup> for Hyola.401 (Table D2). Average seed yield of Clearwater was higher than Sunrise in each of the 6 years tested, albeit that the difference was only significant (P<0.05) in three of the six years.

Clearwater also was compared to the cultivar '45A71' and selection 'PM.2' in yield trials at two locations in 2004 and 2005 where seedlings in the trials were sprayed with 0X, 2X (equivalent to 70 g a.i. ha<sup>-1</sup>), and 3X (equivalent to 105 g a.i. ha<sup>-1</sup>) rates of imazamox herbicide at the 4-5 leaf stage (Table D3). Averaged over years and sites, seed yield of Clearwater was almost identical with 0X herbicide rate (1,748 kg ha<sup>-1</sup>) compared to 2X herbicide rate (1,730 kg ha<sup>-1</sup>). Plant stand counts, plant height, days to flower start and seed oil content of Clearwater remained constant irrespective of imazamox herbicide application rates.

In a similar study conducted in 2004 and 2005 at two locations the performance of Clearwater was recorded when planted into soils previously treated with Pursuit® (Ammonium salt of imazethapyr (±)-2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-5-ethyl-3-pyridinecarboxyic acid) at 0X, 0.5X (equivalent to 17.5 g a.i. ha<sup>-1</sup>), 1X (equivalent to 35 g a.i. ha<sup>-1</sup>)and 2X (equivalent to 70 g a.i. ha<sup>-1</sup>) recommended application rates, and thereafter seedlings sprayed with imazamox at a 0X, 1X (equivalent to 35 g a.i. ha<sup>-1</sup>), and 2X (equivalent to 70 g a.i. ha<sup>-1</sup>) foliar application rate (Table D4). Imazethapyr and imazamox are both imidazoline class herbicides. Seed yield of Clearwater remained constant over all herbicide and non-herbicide treatments, except in Moscow in 2005, again stressing the high tolerance of Clearwater to imazamox herbicide residue in the soil and foliar applications of this herbicide.

Over the 70 year-sites of testing, oil content of Clearwater ranged from a high of 443 g kg<sup>-1</sup> to a low of 361 g kg<sup>-1</sup> (data not shown), and an average oil content of 391 g kg<sup>-1</sup>, which was not significantly different from either Sunrise or Hyola.401 (Table D5). Seed oil fatty acid composition of Clearwater was very similar to Sunrise, where the primary fat was oleic acid. Clearwater seed oil consistently contained less than 7 g kg<sup>-1</sup> erucic acid, less than 80 g kg<sup>-1</sup> linolenic acid and more than 620 g kg<sup>-1</sup> oleic acid (Table D6). Total glucosinolate content of Clearwater seed meal was low (average of 9.1 µmol g<sup>-1</sup> of defatted seed meal) (Table D7). Primary glucosinolate was 2-hydroxy-3-butenyl (average 5.7 µmol g<sup>-1</sup>) followed by 3-butenyl (average 2.9 µmol g<sup>-1</sup>).

Dec. 16, 2013 2:52 PM





**Table D1.** Days to flower, plant height, lodging and seed shatter of Clearwater, Sunrise, and Hyola.401 spring rapeseed tested in regional trials throughout the Pacific Northwest region between 2001 and 2007.

Cultivar	Days to	Plant	Lodging	Seed
	Flower	Height		Shatter
	days	cm	1-9 scale	1-9 scale
	28 year-sites	24 year-sites	12 year-sites	2 year-sites
Clearwater	58.8	127.8	8.3	7.6
Sunrise	58.6	118.9	6.6	7.5
Hyola.401	54.5	106.7	7.7	8.8
Mean	57.3	117.8	7.5	8.0
LSD 5%	1.2	7.86	1.0	0.6
Sig	**	*	**	*

**Table D2.** Seed yield of Clearwater, Sunrise, and Hyola.401 spring rapeseed tested in regional trials throughout the Pacific Northwest region between 2001 and 2007.

	Seed Yield								
Cultivar	Average	Rank	2007	2006	2005	2004	2003	2002	2001
			10 sites	12 sites	11 sites	13 sites	7 sites	12 sites	14 sites
			-			kg ha <sup>-1</sup>			
Clearwater	1,748	2	1,872	1,870	1,739	1,622	1,318	1,649	2,166
Sunrise	1,646	3		1,645	1,679	1,510	1,289	1,630	2,125
Hyola.401	1,986	1	2,176	2,035	1,818	2,129	1,509	1,860	2,373
Mean	1,793		2,024	1,850	1,745	1,754	1,372	1,713	2,221
LSD 5%	313		312	-	-	233	113	-	155
Sig			*	n.s.	n.s.	**	**	n.s.	*

Dec. 16, 2013 2:52 PM

**Table D3.** Seed yield, plant stand counts, Julian days to flower start, plant height after flower and seed oil content of Clearwater, 45A71 and PM.2, grown at three locations after Beyond (imazamox) herbicide was applied at the 0X, 2X and 3X application rates.

	Beyond	Seed	Plant	Flower	Plant	Seed
Identifier	Rate	Yield	Stand	Start	Height	Oil
		- lb/a -	- # m <sup>-1</sup> -	days	cm	%
Clearwater	0X	1,748	43	179	120	35
Clearwater	2X	1,730	47	179	123	35
Clearwater	3X	1,571	50	179	123	34
45A71	0X	1,349	52	178	119	35
45A71	2X	1,581	55	178	121	35
45A71	3X	1,643	55	178	121	35
PM.2	0X	1,166	46	181	124	35
PM.2	2X	866	40	181	119	35
PM.2	3X	866	31	182	122	35
Average		1,391	47	180	121	35
s.e.		185.0	n.s.	0.5	3.4	0.5
LSD 5%		543.0	-	1.4	9.9	1.6

**Table D4.** Seed yield of Clearwater grown at Moscow and Genesee in 2004 and 2005 with four rates of Pursuit herbicide applied pre-plant incorporated (0X, 0.5X, 1X and 2X) and three rates of Beyond applied foliar (0X, 1X, and 2X).

	Pursuit	Beyond	Average	Moscow	Genesee	Moscow	Genesee
Cultivar	Rate	Rate		2005	2005	2004	2004
					kg ha <sup>-1</sup>		
Clearwater	2X	2X	1,427	887	795	2,060	1,967
Clearwater	2X	1X	1,396	1,023	657	1,861	2,044
Clearwater	2X	0X	1,478	820	725	2,091	2,276
Clearwater	1X	2X	1,414	950	692	1,977	2,036
Clearwater	1X	1X	1,374	1,222	530	1,896	1,846
Clearwater	1X	0X	1,387	762	616	2,295	1,875
Clearwater	0.5X	2X	1,368	698	730	2,068	1,976
Clearwater	0.5X	1X	1,394	1,037	628	1,705	2,206
Clearwater	0.5X	0X	1,379	658	746	2,045	2,066
Clearwater	0X	2X	1,269	783	627	1,779	1,889
Clearwater	0X	1X	1,201	841	553	1,465	1,944
Clearwater	0X	0X	1,102	652	634	1,651	1,471
Average			1,349	861	661	1,908	1,966
LSD 5%			n.s.	156	n.s.	n.s.	n.s.

**Table D5.** Seed oil content of Clearwater, Sunrise, and Hyola.401 spring rapeseed tested in regional trials throughout the Pacific Northwest region between 2001 and 2007.

	Oil Content								
Cultivar	Average	Rank	2007	2006 11	2005	2004	2003	2002	2001
			8 sites	sites	11 sites	11 sites - mg kg <sup>-1</sup>	6 sites	10 sites	13 sites
Clearwater	391	1	387	404	379	391	378	387	411
Sunrise	390	2		402	381	387	376	387	407
Hyola.401	383	3	377	397	372	391	358	386	400
Mean	388		382	401	377	390	371	387	406
LSD 5%	70		18	12	13	18	21	16	17
Sig	n.s.		*	n.s.	n.s.	**	**	n.s.	*

**Table D6.** Fatty acid profile of Clearwater and Sunrise spring canola.

		Fatty Acid Profile							
Cultivar	16:0†	18:0	18:1	18:2	18:3	20:1	22:1		
				<mark>y</mark> ng kg	·1				
Clearwater	41	22	621	195	76	14	7		
s.e.	1.4	1.3	6.2	3.0	5.2	1.4	4.5		
Sunrise	50	19	617	205	94	16	4		
s.e.	0.5	3.0	6.2	7.5	5.4	0.3	0.3		

<sup>† 16:0=</sup>Steric acid; 18:0=Palmitic acid; 18:1=Oleic acid; 18:2 = linoleic acid; 18:3 = linolenic acid; 20:1 = eicoseneic acid; 22:1 = erucic acid

**Table D7.** Glucosinolate profile of Clearwater and Sunrise spring canola.

			Glucosino	olate	
Cultivar	Total	Butenyl †	Hy-butenyl	Penteny	Hy-Pentenyl
			umol g	1	
Clearwater	9.135	2.902	5.747	0.477	Tr
s.e.	0.350	0.107	0.231	0.021	
Sunrise	6.163	2.094	3.839	0.261	Tr
s.e.	0.175	0.067	0.082	0.042	

<sup>† 3-</sup>butenyl glucosinolate; 2-hydroxy-3-butenyl glucosinolate; 4-pentenyl glucosinolate; 2-hydroxy-4-pentenyl glucosinolate

REPRODUCE LOCALLY, Include form number and editi	on date on all reproductions.	ORM APPROVED - OMB No. 0581-0055	
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE  EXHIBIT E  STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held confidential until the certificate is issued (7 U.S.C. 2426).		
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME	
Jack Brown University of Idaho	UI.C.1.04	Clearwater	
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)	
PSES, CALS Morrill Hall 414	208 885 7078	208 885 7760	
PO Box 442339 443003 875 Perimeter Drive,	7. PVPO NUMBER		
University of Idaho MS3003			
Moscow, ID 83844- <del>2339</del> -3003	#201100051		
8. Does the applicant own all rights to the variety? Mark an "X" in the appropriate block. If no, please explain.			
9. Is the applicant a U.S. national or a U.S. based entity? If no, give name of country.  X YES NO			
10. Is the applicant the original owner? YES NO If no, please answer one of the following:			
a. If the original rights to variety were owned by individual(s), is (are) the original owner(s) a U.S. National(s)?  YES  NO  If no, give name of country			
b. If the original rights to variety were owned by a company(ies), is (are) the original owner(s) a U.S. based company?  NO If no, give name of country			
11. Additional explanation on ownership (Trace ownership from original breeder to current owner. Use the reverse for extra space if needed):			
11. Additional explanation on ownership (Trace ownership noth ongi	mar brooder to carroin owner. God the		
	*		
	r.		
PLEASE NOTE:			
Plant variety protection can only be afforded to the owners (not licensees) who meet the following criteria:			

- 1. If the rights to the variety are owned by the original breeder, that person must be a U.S. national, national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 2. If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed the final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definitions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, parental status, parental status, perental status, pe

To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

ST470-E (07-09) designed by the Plant Variety Protection Office

REPRODUCE LOCALLY, Include form number and date on all reproductions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The v OMB control number for this information collection is estimated to average 5 minutes per response, including the time for reviewing instructions searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an Individual's income is derived from any public assistance program (Not all prohibited bases apply to programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and

To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

> **U.S. DEPARTMENT OF AGRICULTURE** AGRICULTURAL MARKETING SERVICE **SCIENCE AND TECHNOLOGY ANT VARIETY PROTECTION OFFICE** BELTSVILLE, MD 20705

**EXHIBIT F DECLARATION REGARDING DEPOSIT** 

NAME OF OWNER (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)  PSES, CALS; PO Box 442339-443003  University of Idaho Morill Hall 414	TEMPORARY OR EXPERIMENTAL DESIGNATION UI.C.1.04
Liniversity of Idaho	University of Idahe Morill Hall 414 Moscow, ID 83844- <del>2839</del> -3003	variety name Clearwater
NAME OF OWNER REPRESENTATIVE (S)	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country)	
Jack Brown	PSES, CALS, Po Box 442339 MS3003,	PVPO NUMBER
Karen Stevenson	University of Idaho 875 Perimeter Drive Moscow, ID 83844 2339 (bt: 5/16/2011)	#201100051

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.

2010 MW 10 PH 6:15